

# Applications and Interpretation Higher Level for IBDP Mathematics

## Practice Paper Set 1 – Paper 1 (120 Minutes)

### Question – Answer Book

#### Instructions

1. Attempt **ALL** questions. Write your answers in the spaces provided in this Question - Answer Book.
2. A graphic display calculator is needed.
3. You are suggested to prepare a formula booklet of Applications and Interpretation for IBDP Mathematics when attempting the questions.
4. Supplementary answer sheets and graph papers will be supplied on request.
5. Unless otherwise specified, **ALL** working must be clearly shown.
6. Unless otherwise specified, numerical answers should be either **EXACT** or correct to **3 SIGNIFICANT FIGURES**.
7. The diagrams in this paper are **NOT** necessarily drawn to scale.
8. Information to be read before you start the exam:



	Marker's Use Only	Examiner's Use Only	
Question Number	Marks	Marks	Maximum Mark
1			5
2			5
3			6
4			5
5			6
6			6
7			7
8			6
9			6
10			5
11			8
12			7
13			6
14			7
15			7
16			5
17			7
18			6
<b>Overall</b>			
<b>Paper 1 Total</b>			<b>110</b>










4. The diagram below shows the Voronoi diagram of three restaurants for take-away meals, A, B and C, in a town bounded by the coordinate axes, the lines  $x=8$  and  $y=8$ , where 1 unit represents 1 km.



The straight line  $L$  is the boundary separating the Voronoi cells of B and C. It is given that  $(4, 4)$  is a point on  $L$ .

- (a) (i) Find the gradient of  $L$ .
- (ii) Hence, find the equation of  $L$ , giving the answer in slope-intercept form.

[4]

Kimberly would like to find a restaurant closest to her office to minimize the delivery time of her meal during lunchtime. The position of her office is at  $(7, 2.5)$ .

- (b) State the reason that she is indifferent from choosing the restaurant B and the restaurant C.

[1]











9. In a supermarket, the weights of apples are normally distributed with mean 140 g and standard deviation 9 g, and the weights of oranges are normally distributed with mean 200 g and standard deviation 14 g. Three apples are randomly chosen. Let  $X$  be the total weight of the selected apples.

(a) Write down

(i) the mean of  $X$  ;

(ii) the variance of  $X$  .

[2]

Three apples and seven oranges are randomly chosen. Let  $Y$  be the total weight of the selected fruits.

(b) Write down

(i) the mean of  $Y$  ;

(ii) the standard deviation of  $Y$  .

[2]

(c) Hence, find  $P(Y \geq 1770)$ .

[2]

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---











13. Two surveys are conducted to measure the residents' satisfaction on the services provided by the community centre. A score from 0 to 10 is used in the surveys, where 0 represents absolute dissatisfaction and 10 represents absolute satisfaction. The table below shows the results of the surveys completed by 6 residents:

Resident	A	B	C	D	E	F
Scores from the first survey ( $x$ )	5	7	3	6	8	8
Scores from the second survey ( $y$ )	4	9	5	5	9	9

The manager of the community centre wants to investigate whether the mean scores of the second survey has improved. A paired  $t$ -test is conducted at a 5% significance level. Define  $d = x - y$ .

- (a) (i) Write down the null hypothesis of the test. [2]
- (ii) Write down the alternative hypothesis of the test. [2]
- (b) Find the  $p$ -value. [2]
- (c) State the conclusion of the test with a reason. [2]

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---







15. A quadratic function is given by  $f(x) = ax^2 + bx + c$ . It is given that the complex roots of  $f(x) = 0$  are  $\frac{1}{2} + \frac{1}{4}i$  and  $\frac{1}{2} - \frac{1}{4}i$ .

(a) Write down the values of

(i)  $\left(\frac{1}{2} + \frac{1}{4}i\right) + \left(\frac{1}{2} - \frac{1}{4}i\right)$ ;

(ii)  $\left(\frac{1}{2} + \frac{1}{4}i\right)\left(\frac{1}{2} - \frac{1}{4}i\right)$ .

[2]

(b) Hence, find the expression of  $f(x)$ , giving the answer in terms of  $a$ .

[3]

The graph of  $f(x)$  passes through  $\left(1, \frac{5}{2}\right)$ .

(c) Find the value of  $a$ .

[2]

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---











